

PROFESSIONAL INFORMATION		PROFESSIONELE INLIGTING	
MAXIPREG		MAXIPREG	
SCHEDULING STATUS		SKEDULERINGSTATUS	
S0		S0	
PROPRIETARY NAME AND DOSAGE FORM		HANDELSNAAM EN DOSEERVORM	
MAXIPREG (capsules)		MAXIPREG (kapsules)	
COMPOSITION		SAMESTELLING	
Each sachet contains 4 capsules:-		Elke sakkie bevat 4 kapsules:-	
1 Multivitamin capsule containing:		1 Multivitamien kapsule met:	
Biotin	5 µg	Biotien	5 µg
Calcium	20 mg	Kalsium	20 mg
Choline	5 mg	Cholien	5 mg
Copper	200 µg	Koper	200 µg
Folic acid	500 µg	Foliensuur	500 µg
Inositol	1 mg	Inositol	1 mg
Iodine	140 µg	Jodium	140 µg
Iron	12 mg	Yster	12 mg
Magnesium	500 µg	Magnesium	500 µg
Manganese	5 µg	Mangaan	5 µg
Phosphorous	15 mg	Fosfaat	15 mg
Potassium	500 µg	Kalium	500 µg
Selenium	50 µg	Selenium	50 µg
Vitamin A (Retinyl palmitate)	2666 iu	Vitamien A (Retinielpalmitaat)	2666 ie
Vitamin B1 (Thiamine Nitrate)	3 mg	Vitamien B1 (Thiamen Nitraat)	3 mg
Vitamin B2 (Riboflavin)	2 mg	Vitamien B2 (Riboflavin)	2 mg
Vitamin B3 (Nicotinamide)	20 mg	Vitamien B3 (Nikotienamied)	20 mg
Vitamin B5 (Pantothenic acid)	5 mg	Vitamien B5 (Pantoteensuur)	5 mg
Vitamin B6 (Pyridoxine HCl)	8 mg	Vitamien B6 (Piridoksiensuur HCl)	8 mg
Vitamin B12 (Cyanocobalamin)	2 µg	Vitamien B12 (Cyanocobalamine)	2 µg
Vitamin C (Ascorbic acid)	40 mg	Vitamien C (Ascorbiensuur)	40 mg
Vitamin D3 (Cholecalciferol)	120 iu	Vitamien D3 (Cholekalsiferol)	120 ie
Vitamin E (Alpha-tocopherol)	20 mg	Vitamien E (Alfa tokoferoel)	20 mg
Zinc	4 mg	Sink	4 mg
1 Marine Fish Oil capsule containing:		1 Mariene Vis Olie kapsule bevat:	
Marine Fish Oil	1000 mg	Mariene vis olie	1000 mg
Providing EPA (Eicosapentaenoic acid)	180 mg	Die verskaffing van EPA (Eikosapentaensuur)	180 mg
DHA (Docosahexaenoic acid)	120 mg	DHA (Dokosahexaensuur)	120 mg
1 Lecithin capsule containing:		1 Lesitin kapsule bevat:	
Lecithin	1000 mg	Lesitin	1000 mg
1 white Calcium capsule containing:		1 wit Kalsium kapsule bevat:	
Calcium	600 mg	Kalsium	600 mg
Vitamin D3	400 iu	Vitamien D3	400 ie
The other ingredients are: Green capsule: soja lecithin, carrageenan, glycerol. Yellow, brown and white capsule: bovine gelatin.		Onaktiewe bestanddele: Groen kapsule: soja lesitin, karrageenan, glicerol. Geel, bruin en wit kapsule: bees gelatien.	
Contains no sugar.		Bevat geen suiker.	
PHARMACOLOGICAL CATEGORY		FARMAKOLOGIESE KATEGORIE	
Health supplement. Category D34.12 Multiple substance formulation.		Gesondheidsaanvulling. Kategorie D34.12 Meervoudige bestanddeel formulering.	
PHARMACOLOGICAL ACTION		FARMAKOLOGIESE WERKING	
The vitamins and minerals can assist in fat, carbohydrate and amino acid metabolism, nucleic acid and protein synthesis. They have antioxidant properties which assist in maintaining proper immune function. The body cannot produce omega-3 fatty acids. MAXIPREG provides a source of omega-3. Calcium is necessary for the development and maintenance of bone strength and may reduce the risk of osteoporosis.		Die vitamiene en minerale kan met vette, koolhidrate en aminosuur metabolisme, nukleïensuur en proteiensintese help. MAXIPREG het anti-oksidant eienskappe wat help met die handhawing van n behoorlike immuno funksie. Die liggaam kan nie omega-3 vetsure produuseer nie en MAXIPREG is 'n bron van omega-3. Kalsium is noodsaalik vir die ontwikkeling en instandhouing van beensterke en kan die risiko van osteoporose verminder.	
Pharmacodynamic properties		Farmakodinamiese eienskappe	
Omega-3 fats are thought to be important for normal neural function. There is preliminary evidence that they might improve retinal, visual, learning, and memory function.		Dit word vermoed dat omega-3 belangrik is vir normale neurale funksie. Daar is bewys dat dit dalk retinale, visuele, leer, en geheue-funksie verbeter.	
Lecithin consists of phospholipids which are the chief constituents of cell membranes. Lecithin also contains omega-6 fatty acids and a small amount of omega-3 fatty acids.		Lesitien bestaan uit fosfolipide wat die hoof bestanddeel van selmembrane is.	
Pharmacokinetic properties		Lesitien bevat ook omega-6 vetsure en 'n klein hoeveelheid omega 3-vetsure.	
Following consumption, polyunsaturated fatty acids, such as the omega-3 and omega-6 fatty acids, are incorporated into cell membranes, where they modulate membrane protein function, cellular signaling, and gene expression.		Farmakokinetiese eienskappe	
INDICATIONS		Na inname van poli-onversadigde vetsure, soos omega-3 en omega-6 vetsure wat in die selmembrane opgeneem word, waar hulle membraan proteine funksie, sellulere sein, en die uitdrukking van gene moduleer.	
MAXIPREG is an all-in-one supplement providing vitamins, minerals, phospholipids and omega-3 formulated to provide what the body needs before, during and after pregnancy.		INDIKASIES	
CONTRAINDICATIONS		MAXIPREG is 'n alles-in-een aanvulling wat vitamiene, minerale, fosfolipiede, kalsium en omega-3 verskaf wat gevormuleer is om voedingsbehoeftes voor, tydens en na swangerskap te voorsien.	
MAXIPREG capsules are contraindicated in patients with known hypersensitivity to any of the ingredients.		MAXIPREG kan ondersteuning bied vir moegheid en die verbetering van konsentrasie.	
WARNINGS AND SPECIAL PRECAUTIONS		KONTRA-INDIKASIES	
Not to be taken by persons allergic to fish. Consult a doctor before taking this product if you are on any other Vitamin A medication. Consult a doctor if you are on anticoagulants.		MAXIPREG is teenaangedui in pasiente wat hipersensitief is vir enige van die bestanddele.	
Special precautions		WAARSKUWINGS EN SPEISALE VOORSORGAATREELS	
Lesitin can cause allergic skin reactions in people with egg and soy allergies.		Moet nie geneem word deur persone wat allergies is vir vis nie. As jy op enige ander vitamiene A medikasie is, raadpleeg jou dokter voordat jy hierdie produk gebruik. Raadpleeg jou dokter as jy op antistollings medisyne is.	
MAXIPREG should not be taken by patients suffering from kidney failure, kidney stones or calcium deposits in the kidneys. MAXIPREG should not be taken by patients suffering from Sarcoidosis as it contains calcium and vitamin D3.		Spesiale voorsorgmaatreels	
Effects on ability to drive vehicles and use machinery: No known effect.		Lesitien kan allergiese vel reaksies veroorsaak in mense met eier en soja allergieë.	
INTERACTIONS		MAXIPREG moet nie geneem word deur pasiënte wat aan niersverstoring, niersteen of kalsiumafsettings in die niere ly nie. MAXIPREG moet nie geneem word deur pasiënte wat aan sarkoidose ly nie, aangesien dit kalsium en vitamien D3 bevat.	
Vitamin A may interfere with Warfarin, reducing the anticoagulant effect.		Efek op die vermoë om motor te bestuur en masjinerie te gebruik: Geen bekende effek.	
Thiazide diuretics may increase calcium blood levels.		INTERAKSIES	
Oral steroids may increase calcium blood levels.		Vitamien A kan met Warfarin inmeng en die vermindering van die antistollingsuitwerking beïnvloed.	
Cardiac glycosides may exhibit greater side effects with calcium supplementation.		Tiasiediureтика kan kalsiumbloede/vlake verhoog.	
Absorption of tetracyclines, fluoroquinolones and bisphosphonates may be inhibited by calcium and should be taken 2 hours before or 4 hours after calcium intake.		Orale sterolede kan kalsiumbloede/vlake verhoog.	
PREGNANCY AND BREASTFEEDING		Hartglikoside kan groter newe-effekte met kalsiumaanvulling.	
Speak to your doctor if you are pregnant or breastfeeding or plan to become pregnant soon.		Absorpsie van tetrasikliene, fluoroquinolone en bisfosfonate kan deur kalsium geïnhieber word en moet 2 uur voor of 4 uur na kalsiuminnname geneem word.	
Taking too much vitamin A can be harmful to your baby.		SWANGER EN BORSVOED	
MAXIPREG can be taken during pregnancy and lactation.		Raadpleeg jou dokter as jy swanger is of borsvoed, of as jy beplan om binnekort swanger te word.	
As calcium and vitamin D3 pass into breast milk, care must be taken if the infant is taking any other supplements containing calcium and vitamin D3.		Te veel vitamiene A kan skadelik vir jou baby wees.	
DOSAGE AND DIRECTIONS FOR USE		MAXIPREG kan tydens swangerskap en laktasie geneem word.	
Persons 18 years and older: 1 sachet with the 4 capsules daily with a meal and half a glass of water.		Aangesien kalsium en vitamien D3 in borsmelk orgaan, wees dus versigtig as die baba enige ander aanvullings wat kalsium en vitamien D3 bevat, neem.	
SIDE EFFECTS		DOSIS EN GEBRUIKSAANWYSINGS	
Nervous system disorders		Persone 18 jaar en ouer: 1 sakkie met die 4 kapsules per dag met ete en 'n halwe glas water.	
Less frequent		NEWE-EFFEKTE	
Gastrointestinal disorders		Senusisteemversteurings	
Less frequent		Minder gereeld	
Skin		Gastrointestinale versteurings	
Less frequent		Minder gereeld	
KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT		Hardlywighed, winderigheid, viserige winde opbrek, scooterbrand, slechte spysvertering en naарheid, helder geel verkleuring van die urine en halitose	
Treatment should be symptomatic and supportive.		Vel	
IDENTIFICATION		Minder gereeld	
Each sachet contains:		Velutslag	
1 green capsule providing vitamins and minerals.		BEKENDE SIMPTOME VAN OÖRDOSERING EN BESONDERHEDE VIR DIE BEHANDELING DAARVAN	
1 clear, light-yellow capsule providing marine fish oil.		Behandeling is simptomaties en ondersteunend.	
1 brown capsule providing lecithin.		IDENTIFIKASIE	
1 white capsule providing calcium and vitamin D3.		Elke sakkie bevat:	
Note that the colour of the lecithin may vary from batch to batch. This is quite normal and is due to varying environmental factors upon harvesting of the oil.		1 groen kapsule met vitamiene en minerale.	
PRESENTATION		1 deursigte, liggeel kapsule met mariene vis olie.	
Each sachet of MAXIPREG contains 4 capsules. There are 28 sachets in a box.		1 bruin kapsule met lesitin.	
STORAGE INSTRUCTIONS		1 wit kapsule met kalsium en vitamien D3.	
Store at or below 25 °C		Let daarop dat die kleur van die Lesitien mag verskil van bondel tot bondel. Dit is heel normaal as gevolg van omgewings faktore met die oes van die olie.	
Protect from light.		AANBIEDING	
Do not use after the expiry date printed on the box.		MAXIPREG bevat 4 kapsules per sakkie. 28 sakkies in 'n boks.	
STORE OUT OF REACH OF CHILDREN		BERGINGSINSTRUKSIES	
REGISTRATION NUMBER		Stoor teen of onder 25 °C.	
Will be allocated by Council upon registration.		Beskerm teen lig.	
NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION		Moet nie na die vervaldatum op die boks gebruik nie.	
Keynote Health (Pty) Ltd		STOOR BIJSTE BEREIK VAN KINDERS.	
Unit 2		REGISTRASIONOMMER	
Penny Lane Park		Sal toegewe word deur die Raad na afloop van registrasie.	
64 Ebonyfield Ave		NAAM EN BESIGHEIDSADRES VAN DIE HOOUER VAN DIE REGISTRASIESERTIFIKAAT	
Springfield Park		Keynote Health (Pty) Ltd	
4091		Unit 2	
DATE OF PUBLICATION		Penny Lane Park	
March 2023		64 Ebonyfield Ave	
This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.		Springfield Park	
		4091	

PATIENT INFORMATION LEAFLET MAXIPREG	PASIËNTINLIGTINGSBLAADJIE MAXIPREG
SCHEDULING STATUS S0	SKEDULERINGSTATUS S0
PROPRIETARY NAME AND DOSAGE FORM MAXIPREG (capsules)	HANDELSNAAM EN DOSEERVORM MAXIPREG (kapsules)
Read all of this leaflet carefully because it contains important information for you. MAXIPREG is available without a doctor's prescription. Nevertheless you still need to use MAXIPREG carefully to get the best results from it.	Lees die hele voubiljet noukeurig deur, want dit bevat belangrike inligting vir jou. MAXIPREG is beskikbaar sonder 'n voorskrif. Nogtans moet jy die gebruiksaanwysings vir MAXIPREG noukeurig volg om die beste resultate te verkry.
• Keep this leaflet. You may need to read it again. • Do not share MAXIPREG with any other person. • Ask your pharmacist if you need more information or advice. • You must see a doctor if your symptoms worsen or do not improve.	• Hou hierdie voubiljet. Jy mag dit dalk weer moet lees. • Moet nie MAXIPREG met enige ander persoon deel nie. • Vra jou apoteker indien jy meer inligting of advies nodig het. • Raadpleeg jou dokter as jou simptome vererger of nie verbeter nie.
WHAT MAXIPREG CONTAINS	WAT MAXIPREG BEVAT
Each sachet contains 4 capsules:- 1 Multivitamin capsule containing:	Elke sakkie bevat 4 kapsules:- 1 Multivitamien kapsule met:
Biotin 5 µg Calcium 20 mg Choline 5 mg Copper 200 µg Folic acid 500 µg Inositol 1 mg Iodine 140 µg Iron 12 mg Magnesium 500 µg Manganese 5 µg Phosphorous 15 mg Potassium 500 µg Selenium 50 µg Vitamin A (Retinyl palmitate) 2666 IU Vitamin B1 (Thiamine Nitrate) 3 mg Vitamin B2 (Riboflavin) 2 mg Vitamin B3 (Nicotinamide) 20 mg Vitamin B5 (Pantothenic acid) 5 mg Vitamin B6 (Pyridoxine HCl) 8 mg Vitamin B12 (Cyanocobalamin) 2 µg Vitamin C (Ascorbic acid) 40 mg Vitamin D3 (Cholecalciferol) 120 IU Vitamin E (Alpha-tocopherol) 20 mg Zinc 4 mg	Biotin 5 µg Kalsium 20 mg Cholien 5 mg Koper 200 µg Folienuur 500 µg Inositol 1 mg Jodium 140 µg Yster 12 mg Magnesium 500 µg Mangaan 5 µg Fosfaat 15 mg Kalium 500 µg Selenium 50 µg Vitamien A (Retinylpalmitaat) 2666 ie Vitamien B1 (Thiamine Nitraat) 3 mg Vitamien B2 (Riboflavin) 2 mg Vitamien B3 (Nikotienamied) 20 mg Vitamien B5 (Pantoteensuur) 5 mg Vitamien B6 (Piridoksiens HC1) 8 mg Vitamien B12 (Cyanocabalamin) 2 µg Vitamien C (Askorbiensuur) 40 mg Vitamien D3 (Cholecalciferol) 120 ie Vitamien E (Alfa tokofeler) 20 mg Sink 4 mg
1 Marine Fish Oil capsule containing: Marine Fish Oil 1000 mg Providing EPA (Eicosapentaenoic acid) 180 mg DHA (Docosahexaenoic acid) 120 mg	1. Marine Vis Olie kapsule bevat: Mariene vis olie 1000 mg Die verskaffing van EPA (Eikospentaensuur) 180 mg DHA (Dokosahexaensuur) 120 mg
1 Lecithin capsule containing: Lecithin 1000 mg	1 Lesitien kapsule bevat: Lesitien 1000 mg
1 white Calcium capsule containing: Calcium 600 mg Vitamin D3 400 IU	1 wit Kalsium kapsule bevat: Kalsium 600 mg Vitamien D3 400 ie
The other ingredients are: Green capsule: soya lecithin, carrageenan, glycerol. Yellow, brown and white capsule: bovine gelatin. Contains no sugar.	Onaktiewe bestanddele: Groen kapsule: soja lesitien, karrageenan glicerol. Geel, bruin en wit kapsule: bees gelatin. Bevat geen suiker.
PHARMACOLOGICAL CATEGORY Health supplement. Category D34.12 Multiple substance formulation.	FARMAKOLOOGIESE KATEGORIE Gesondheidsaamvulling. Kategorie D34.12 Meervoudige bestanddeel formulering.
WHAT MAXIPREG IS USED FOR Calcium is necessary for the development and maintenance of bone strength and may reduce the risk of osteoporosis. MAXIPREG is an all-in-one supplement providing vitamins, minerals, phospholipids and omega-3, formulated to provide what the body needs before, during and after pregnancy.	WAARVOOR MAXIPREG GEBRUIK WORD MAXIPREG is 'n alles-in-een aanvulling wat vitamiene, minerale, fosfolipide, kalsium en omega-3, verskaf wat geformuleer is om voedingsbehoefte voor, tydens en na swangerskap te voorsien.
BEFORE YOU TAKE MAXIPREG: Do not take MAXIPREG • If you are hypersensitive (allergic) to any of the ingredients. Tell your doctor if you are not sure about any of the above. Take special care with MAXIPREG MAXIPREG should not be taken by patients suffering from kidney failure, kidney stones or calcium deposits in the kidneys. MAXIPREG should not be taken by patients suffering from Sarcoidosis as it contains calcium and vitamin D3. Not to be taken by persons allergic to fish. Consult a doctor before taking this product if you are on any other Vitamin A medication. MAXIPREG can cause allergic skin reactions in people with egg and soy allergies. Taking MAXIPREG with food and drink: MAXIPREG should be taken with water and a meal.	VOORDAT JY MAXIPREG NEEM: Moet nie MAXIPREG gebruik: • Indien jy hypersensitief (allergie) is vir enige van die bestanddele. Raadpleeg jou dokter indien jy nie seker is oor enige van bogenoemde nie. Neem spesiale voorvars met MAXIPREG MAXIPREG moet nie geneem word deur pasiënte wat aan niersversaking, nierstene of kalsiumafsettings in die niere ly nie. MAXIPREG moet nie geneem word deur pasiënte wat aan sarkoïdose ly nie, aangesien dit kalsium en vitamien D3 bevat. Moet nie geneem word deur persone wat allergies vir vis is nie. Raadpleeg jou dokter voordat jy hierdie produk gebruik as jy op enige ander vitamiene A medikasie is. MAXIPREG kan allergiese vel reakções in mense met eier en soja allergie veroorsaak. Die neem van MAXIPREG met kos en drank: MAXIPREG moet met water en 'n maaltyd geneem word. SWANGERSKAP EN BORSVOED As jy swanger is of jou baby borsvoed, raadpleeg jou dokter, apoteker of 'n ander professionele gesondheidsorg werker vir advies voordat jy hierdie medisyne neem. MAXIPREG kan tydens swangerskap en laktasie geneem word. Aangesien kalsium en vitamien D3 in borsmilk orgaan, wees dus versigtig as die baba enige ander aanvullings wat kalsium en vitamien D3 bevat, neem.
PREGNANCY AND BREASTFEEDING If you are pregnant or breastfeeding your baby, please consult your doctor, pharmacist or other healthcare professional for advice before taking this medicine. MAXIPREG can be taken during pregnancy and lactation. As calcium and vitamin D3 pass into breast milk, care must be taken if the infant is taking any other supplements containing calcium and vitamin D3. Effects on ability to drive vehicles and use machinery: No known effect. Taking other medicine with MAXIPREG : Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines.) Tell your doctor if you are taking any anticoagulants such as Warfarin. Thiazide diuretics may increase calcium blood levels. Oral steroids may increase calcium blood levels. Cardiac glycosides may exhibit greater side effects with calcium supplementation. Absorption of tetracyclines, fluoroquinolones and bisphosphonates may be inhibited by calcium and should be taken 2 hours before or 4 hours after calcium intake.	Efek op die vermoë om motor te bestuur en masjinerie te gebruik: Geen bekende effek. Gebruik van ander medisyne saam met MAXIPREG Vertel jou gesondheidsorg professioneel as jy enige ander medisyne gebruik. (insluitend aanvullende van tradisionele medisyne.) Sorg dat jou dokter weet indien jy enige ander antistolmiedels soos Warfarin neem. Raadpleeg jou dokter of apoteker as jy enige van die volgende medisyne gebruik: Tiasiediureтика kan kalsiumblouvlakte verhoog. Orale steroidiese kan kalsiumblouvlakte verhoog. Hartglikosiede kan groter newe-effekte met kalsiumaamvulling toon. Tetrasikline, fluorokinolone en bisfosfonate. Absorpsie van hierdie medisyne kan deur kalsium geinhibit word en moet 2 uur voor of 4 uur na kalsiumaamvulling geneem word.
HOW TO TAKE MAXIPREG Do not share medicines with any other person. Always take MAXIPREG exactly as indicated or as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure. The dosage is: Persons 18years and older: one sachet with the 4 capsules daily with a meal and half a glass of water. Do not take more than the recommended dose. If you take more MAXIPREG than you should In the event of overdose, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre. If you forgot to take / missed a dose of MAXIPREG Take your missed dose when you remember. Do not take a double dose to make up for forgotten individual doses. Effects when treatment with MAXIPREG is stopped If taking the dose as indicated, then there should be no adverse effects when stopping use of this medication.	HOE OM MAXIPREG TE NEEM Moet nie jou medisyne met enige ander persone deel nie. Neem MAXIPREG presies soos aangedui of soos deur jou dokter voorgeskryf. Raadpleeg jou dokter of apoteker indien jy onseker is. Die normale dosis is: Persone 18 jaar en ouer: een sakkie met die 4 kapsules per dag met ete en 'n halwe glas water. Moet nie die vastgestelde dosis onskryf nie. As jy meer MAXIPREG neem as wat jy behoort In die geval van oordopsering, raadpleeg jou dokter of apoteker. Indien hulle nie beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum. As jy vergeet / negelat het om 'n dosis van MAXIPREG te neem Neem die oorgeseelde dosis sodra jy onthou daarvan. Moet nie 'n dubbele dosis neem om op te maak vir individuele dosisse wat jy oorgeseel het nie. Uitwerking na behandelung met MAXIPREG gestaak: Indien die dosis soos aangedui geneem word, sal daar geen newe-effekte wees wanneer die gebruik van MAXIPREG gestaak word nie.
POSSIBLE SIDE EFFECTS MAXIPREG has a few uncommon side effects such as headache, sleep disturbances, constipation, flatulence, heartburn, indigestion and nausea; bright yellow discolouration of the urine, halitosis, 'fishy' burp and rash. Not all side effects reported for MAXIPREG are included in this leaflet. Should your general health worsen or if you experience any untoward effects while taking this medicine, please consult your doctor, pharmacist or other health care professional for advice. If any of the following happens, stop taking MAXIPREG and tell your doctor immediately or go to the casualty department at your nearest hospital: • Spontaneous bleeding. • Diarrhoea, nausea, vomiting, lack of muscle tone, and weakness. • Increase in blood glucose levels in diabetic patients. If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.	MOONLIKE NEWE-EFFEKTE MAXIPREG het 'n paar ongewone newe-effekte soos hoofpyn, slaapversteurings, hardlywigheid, winderigheid, soolbrand, slechte spysvertering en naerheid, helder geel verkleuring van die urine, halitose en die opbrek van viserige windie. Nie alle newe-effekt aangemeeld vir MAXIPREG is in hierdie pamphlet ingesluit nie. Indien jou algemene gesondheid verswak of as jy 'n ongewenste effekervaar, raadpleeg jou dokter, apoteker of 'n ander professionele gesondheidsorg werker vir advies. Indien enige van die volgende gebeur, staak gebruik van MAXIPREG en raadpleeg jou dokter onmiddellik, of gaan na die ongevalle-adfeling van jou naaste hospitaal: • Spontane bloeding. • Diarree, naerheid, braking, 'n gebrek aan spiertonus, en swakheid. • 'n Toename in bloedglukoselvlakte in diabetiese pasiente. As jy enige newe-effekte ondervind wat nie in hierdie pamphlet genoem word nie, laat asseblief jou dokter of apoteker weet.
STORING AND DISPOSING OF MAXIPREG Store at or below 25 °C Protect from light and moisture. Do not store in a bathroom. Do not use after the expiry date stated on the carton. Return all unused medicine to your pharmacist. Do not dispose of unused medicine in drains or sewage systems (e.g. toilets).	HOE OM MAXIPREG TE STOOR EN DAARVAN ONTSLAE TE RAAK Stoor teen of onder 25 °C. Die kapsules moet in lig en vog bestande hours gestoor word. Moet nie in 'n badkamer stoor nie. Moet nie dae verwoldaplastuur op die karton gebruik nie. Neem alle ongebruikte medisyne terug na jou apoteker. Moet nie ongebruikte medisyne in dreine of rooiloktelsels (bv. toilette) weggooi nie. STOOR ALLE MEDISyne BIJTE BEREIK VAN KINDERS.
PRESENTATION OF MAXIPREG Each sachet of MAXIPREG contains 4 capsules. There are 28 sachets in a box.	AANBIEDING VAN MAXIPREG MAXIPREG bevat 4 kapsules per sakkie, 28 sakkies in 'n boks
IDENTIFICATION OF MAXIPREG Each sachet contains: 1 green capsule providing vitamins and minerals. 1 clear, light - yellow capsule providing marine fish oil. 1 brown capsule providing lecithin. 1 white capsule providing calcium and vitamin D3. Note that the colour of the lecithin may vary from batch to batch. This is quite normal and is due to varying environmental factors upon harvesting of the oil.	IDENTIFIKASIE VAN MAXIPREG Elke sakkie bevat: 1 groen kapsule met vitamiene en minerale. 1 deursigtige liggeel kapsule met mariene vis olie. 1 bruin kapsule met lecithien. 1 wit kapsule met kalsium en vitamien D3. Let daarop dat die kleur van die Lesitien mag verskil van bondel tot bondel. Dit is heel normaal as gevolg van omgewings faktore met die oes van die olie.
REGISTRATION NUMBER Will be allocated by Council upon registration.	REGISTRASIONOMMER Sal toegewe word deur die Raad na afloop van registrasie.
NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION Keynote Health (Pty) Ltd Unit 2 Penny Lane Park 64 Ebonyfield Ave Springfield Park 4091	NAAM EN BESIGHEIDSADRES VAN DIE HOUER VAN DIE REGISTRASIESERTIFIKAAT Keynote Health (Pty) Ltd Unit 2 Penny Lane Park 64 Ebonyfield Ave Springfield Park 4091
DATE OF PUBLICATION March 2023	DATUM VAN PUBLIKASIE VAN DIE VOUBLILJET Maart 2023
This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.	Hierdie ongeregistreerde medisyne was nie geëvalueer deur die SAHPRA vir kwaliteit, veiligheid of gebruiksin tensies.