

PROFESSIONAL INFORMATION FLEXOR CAPSULES	PROFESSIONELE INLIGTING FLEXOR KAPSULES	
SCHEDULING STATUS SO	SKEDULERINGSTATUS SO	
PROPRIETARY NAME AND DOSAGE FORM FLEXOR (capsules)	HANDELSNAAM EN DOSEERVORM FLEXOR (kapsules)	
COMPOSITION Each capsule contains: Glucosamine Sulphate 375 mg Chondroitin Sulphate 125 mg Collagen Type II 75 mg Vitamin D3 (cholecalciferol) 250 iu MSM (methylsulphonylmethane) 200 mg Papain 63 mg The other ingredients are: Di-calcium phosphate, magnesium stearate, microcrystalline cellulose, piroxil. Shell ingredients: Bovine gelatine, colourants, iron oxide, titanium dioxide. Contains no sugar or artificial sweeteners.	SAMESTELLING Elke kapsule bevat: Glucosamine sulfaat 375 mg Chondroitin sulfaat 125 mg Kollageen Tipe II 75 mg Vitamiën D3 (cholekalsiferol) 250 ie MSM (methylsulphonylmethane) 200 mg Papaïen 63 mg Die ander bestanddele is: Di-calcium phosphate, magnesium stearate, microcrystalline cellulose, piroxil. Dop: Bees gelatien, kleurstowwe, ysteroksied, titaandioksied. Bevat geen suiker of kunsmatige versoeters.	
PHARMACOLOGICAL CATEGORY Complementary medicine - Health supplement. D34.12 Multiple substance formulation.	FARMAKOLOGIESE KATEGORIE Komplementêre medisyne - Gesondheidsaanvulling. D34.12 Meervoudige bestanddeel formulering.	
PHARMACOLOGICAL ACTION Flexor Capsules provide nutrients that contribute to the body's natural healing processes in support of joint mobility.	FARMAKOLOGIESE WERKING Flexor Kapsules verskaf voedingstowwe wat bydra tot die liggaam se natuurlike genesings prosesse ter ondersteuning van gewrigs mobiliteit.	
INDICATIONS FLEXOR CAPSULES contain a combination of nutrients specifically formulated to support healthy joint function.	INDIKASIES FLEXOR KAPSULES bevat 'n kombinasie van voedingstowwe wat spesifiek geformuleer is om gesonde gewrigsfunksie te ondersteun.	
CONTRAINDICATIONS Hypersensitivity to any of the ingredients, including excipients. Fish or shellfish allergy, hypercalcaemia.	KONTRA-INDIKASIES Hipersensitiwiteit vir enige van die bestanddele, insluitend aanvullings, vis of skulpvis allergie en hiperkalsemie.	
WARNINGS AND SPECIAL PRECAUTIONS • Flexor Capsules may increase the anti-coagulant effect of Warfarin or other anti-coagulants. (See INTERACTIONS). • The safety of this preparation in pregnancy and lactation has not been established (See PREGNANCY AND LACTATION). Not suitable for children.	WAARSKUWINGS EN SPESIALE VOORSORGMATREELS • FLEXOR KAPSULES kan die anti-stollings effek van Warfarin en ander anti-stollings middels verhoog. (SIEN INTERAKSIES) • Die veiligheid van die preparaat in swangerskap en laktasie is nie vasgestel nie. (SIEN SWANGERSKAP EN LAKTASIE). Nie geskik vir kinders.	
INTERACTIONS FLEXOR CAPSULES may increase the anti-coagulant effect of Warfarin or other anti-coagulants. (See WARNINGS AND SPECIAL PRECAUTIONS).	INTERAKSIES FLEXOR KAPSULES mag die anti-stollings effek van middels soos Warfarin en ander anti-stollings middels verhoog. (SIEN WAARSKUWINGS EN SPESIALE VOORSORG)	
PREGNANCY AND LACTATION The safety of FLEXOR CAPSULES in pregnancy and lactation has not been established. Women of Childbearing Potential: Should you be planning to fall pregnant, speak to your doctor before using this preparation.	SWANGERSKAP EN LAKTASIE Die veiligheid van FLEXOR KAPSULES in swangerskap en laktasie is nie vasgestel nie. Vroue met die potensiaal om kinders te baar: As jy van plan is om swanger te word, raadpleeg eers jou dokter voordat jy die preparaat gebruik.	
DOSAGE AND DIRECTIONS FOR USE Adults over 18 years: take 2-4 capsules daily. Maximum 4 capsules per day. Dose may be taken as a single dose or divided into two doses. Take 4 capsules daily for severe joint damage or 2 capsules daily for joint maintenance.	DOSIS EN GEBRUIKSAANWYSINGS Volwassenes bo 18 jaar: neem 2-4 kapsules daagliks. Maksimum 4 kapsules per dag. Die dosis mag as 'n enkele dosis geneem, of in twee dosisse verdeel word. Neem 4 kapsules daagliks vir ernstige gewrigskade of 2 kapsules daagliks vir gewrigsonderhoud.	
SIDE EFFECTS	NEWE-EFFEKTE	
Nervous system disorders: Less frequent	Headache	Senusistiemversteurings: Minder gereeld
Cardiac disorders: Less frequent	Extrasystoles	Kardiale versteurings: Minder gereeld
Gastrointestinal disorders: Frequent	Nausea, heartburn, diarrhea, constipation, bloating, dyspepsia and belching.	Gastrointestinale versteurings: Gereelde
Skin and subcutaneous tissue disorders: Frequent Less frequent	Allergic skin reactions, urticaria, rash and pruritus. Eyelid edema.	Vel en subkutane weefsel: Gereelde Minder gereeld
		Hoofpyn
		Ekstrasistolle
		Naarheid, sooi-brand, diarree, hardlywigheid, opgeblaasheid, dispepsie en winde opbreek.
		Allergiese vel reaksies, urtikarie, uitslag en jeuk. Ooglid edeem.
KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT Treatment should be symptomatic and supportive. (see side effects).	BEKENDE SIMPTOME VAN OORDOSERING EN BESONDERHEDE VIR DIE BEHANDELING DAARVAN Behandeling moet simptome en ondersteunend wees. (SIEN NEWE EFFEKTE).	
IDENTIFICATION Off-white, 2 part capsule.	IDENTIFIKASIE Afwit, 2 deel kapsule.	
PRESENTATION Packs of 60 capsules or 120 capsules.	AANBIEDING Pakke van 60 kapsules of 120 kapsules.	
STORAGE INSTRUCTIONS Store at or below 25 °C. Protect from light and store in a cool, dry place. Keep in original packaging. Do not use after the expiry date printed on the bottle or box. STORE OUT OF REACH OF CHILDREN.	BERGINGSINSTRUKSIES Hou teen of onder 25 °C. Beskerm teen lig en stoor in 'n koel, droë plek. Stoor in oorspronklike verpakking. Moet nie na die vervaldatum op die bottel of karton gebruik nie. STOOR BUITE BEREIK VAN KINDERS.	
REGISTRATION NUMBER Will be allocated by Council upon registration.	REGISTRASIENOMMER Sal toegeken word deur die Raad na afloop van registrasie.	
NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION Keynote Health (Pty) Ltd 6 Heron Park 80 Corobrik Road Riverhorse Valley Durban 4017	NAAM EN BESIGHEIDSAADRES VAN DIE HOUER VAN DIE REGISTRASIESERTIFIKAAT Keynote Health (Pty) Ltd 6 Heron Park 80 Corobrik Road Riverhorse Valley Durban 4017	
DATE OF PUBLICATION March 2022	DATUM VAN PUBLIKASIE Maart 2022	
This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.	Hierdie ongeregistreerde medisyne was nie geëvalueer deur die SAHPRA vir kwaliteit, veiligheid of gebruik intensies.	

PATIENT INFORMATION LEAFLET FLEXOR CAPSULES	PASÏËNTINLIGTINGSBLAADJIE FLEXOR KAPSULES
SCHEDULING STATUS SO	SKEDULERINGSSTATUS SO
PROPRIETARY NAME AND DOSAGE FORM FLEXOR CAPSULES (capsules)	HANDELSNAAM EN DOSEERVORM FLEXOR (kapsules)
PHARMACOLOGICAL CATEGORY Complementary medicine - Health supplement. D34.12 Multiple substance formulation.	FARMAKOLOGIESE KATEGORIE Komplementêre medisyne - Gesondheidsaanvulling. D34.12 Meervoudige bestanddeel formulering.
Read all of this leaflet carefully because it contains important information for you. • Keep this leaflet. You may need to read it again. • Do not share FLEXOR CAPSULES with any other person. • Ask your pharmacist if you need more information or advice. • You must see a doctor if your symptoms worsen or do not improve.	Lees die hele voubiljet noukeurig deur, want dit bevat belangrike inligting vir jou. • Hou hierdie voubiljet. Jy mag dit dalk weer moet lees. • Moet nie FLEXOR KAPSULES met enige ander persoon deel nie. • Vra jou apteker indien jy meer inligting of advies nodig het. • Raadpleeg jou dokter as jou simptome vererger of nie verbeter nie.
WHAT FLEXOR CAPSULES CONTAIN The active substances are: Glucosamine Sulphate 375 mg Chondroitin Sulphate 125 mg Collagen Type II 75 mg Vitamin D3 (cholecalciferol) 250 iu MSM (methylsulphonylmethane) 200 mg Papain 63 mg The other ingredients are: Di-calcium phosphate, magnesium stearate, microcrystalline cellulose, piroxil. Shell ingredients: Bovine gelatine, colourants, iron oxide, titanium dioxide. Contains no sugar or artificial sweeteners.	WAT FLEXOR KAPSULES BEVAT Die aktiewe bestanddele is: Glucosamine sulfaat 375 mg Chondroitin sulfaat 125 mg Kollageen Tipe II 75 mg Vitamiën D3 (cholecalciferol) 250 iu MSM (methylsulphonylmethane) 200 mg Papain 63 mg Die ander bestanddele is: Di-calcium phosphate, magnesium stearate, microcrystalline cellulose, piroxil. Dop: Bees gelatien, kleurstowwe, ysteroksied, titaandioksied. Bevat geen suiker of kunsmatige versoeters.
WHAT FLEXOR CAPSULES ARE USED FOR: FLEXOR CAPSULES contain a combination of nutrients specifically formulated for healthy joint function.	WAARVOOR FLEXOR KAPSULES GEBRUIK WORD FLEXOR KAPSULES bevat 'n kombinasie van voedingstopwe spesifiek geformuleer vir gesonde gewrigs funksie.
BEFORE YOU TAKE FLEXOR CAPSULES Do not take FLEXOR CAPSULES: • If you are hypersensitive (allergic) to the active substance of any of the ingredients. • If you are allergic to fish or shellfish. • If you have hypercalcaemia (elevated levels of calcium in the blood). • If you are pregnant or breastfeeding. Take special care with FLEXOR CAPSULES • If you are on anticoagulant therapy (blood thinning agents) (also see Taking other medicines with Flexor), consult your doctor first. Pregnancy and Breastfeeding: The safety of this preparation during pregnancy and breastfeeding has not been established. Women of Childbearing Potential: Should you be planning to fall pregnant, speak to your doctor before using this preparation. Driving and using machinery: Effects not known. Taking other medicines with FLEXOR CAPSULES Always tell your healthcare professional if you are taking any other medicine. (This includes complementary or traditional medicines.) Tell your doctor if you are taking any other medicine such as Warfarin. Flexor may increase the anticoagulant effect of Warfarin. Flexor is not suitable for children.	VOORDAT JY FLEXOR KAPSULES NEEM Moet nie FLEXOR KAPSULES gebruik: • Indien jy hipersensitief (allergies) is vir enige van die bestanddele. • Indien jy allergies is vir vis of skulpvis. • Indien jy hiperkalemie het. (Hoë vlakke van kalsium in die bloed). • Indien jy swanger is of borsvoed. Neem spesiale voorsorg met FLEXOR KAPSULES Indien jy op antistollings terapie medikasie (bloed verdunnings medikasie.) (sien ook Gebruik van ander medikasie saam met FLEXOR KAPSULES), raadpleeg eers jou dokter. Swangerskap en Borsvoeding Die veiligheid van die preparaat gedurende swangerskap en borsvoeding is nog nie vasgestel nie. Motor bestuur en gebruik van masjinerie: Effek onbekend. Gebruik van ander medisyne saam met FLEXOR KAPSULES Vertel jou gesondheidsorg professionele as jy enige ander medikasie gebruik. (insluitend aanvullende of tradisionele medisyne) Raadpleeg jou dokter as jy enige ander medikasie soos Warfarin neem. Flexor mag die uitwerking van antistolterapie medikasie soos Warfarin verhoog. Flexor is nie geskik vir kinders.
HOW TO TAKE FLEXOR CAPSULES Do not share medicines with any other person. Always take FLEXOR CAPSULES exactly as indicated or as your doctor has instructed you. You should check with your doctor or pharmacist if you are unsure. Adults over 18 years take 2-4 capsules daily. Maximum 4 capsules per day. Dose may be taken as a single daily dose or divided into two doses. Take 4 capsules daily for severe joint damage or 2 capsules daily for joint maintenance. If you take more FLEXOR CAPSULES than you should In the event of overdosage, consult your doctor or pharmacist. If neither is available, contact the nearest hospital or poison control centre. If you forget to take / missed a dose of FLEXOR CAPSULES Take your missed dose when you remember. Do not take a double dose to make up for forgotten individual doses. Effects when treatment with FLEXOR CAPSULES is stopped If taking the dose as indicated, then there should be no adverse effects when stopping use of this medication.	HOE OM FLEXOR KAPSULES TE NEEM Moet nie jou medisyne met enige ander persoon deel nie. Neem altyd FLEXOR KAPSULES presies soos aangedui of soos jou dokter voorgeskryf. Raadpleeg jou dokter of apteker indien jy onseker is. Volwasse nes bo 18 jaar: 2-4 kapsules daaglik. Maksimum dosis is 4 kapsules daaglik. Die daaglikse dosis kan as 'n enkele dosis of in twee dosisse verdeel word. Neem 4 kapsules daaglik vir ernstige gewrigskade of 2 kapsules daaglik vir gewrigsonderhoud. As jy meer FLEXOR KAPSULES neem as wat jy behoort In die geval van oordosering, raadpleeg jou dokter of apteker. Indien hulle nie beskikbaar is nie, kontak die naaste hospitaal of gifbeheersentrum. As jy vergeet/nagelaat het om 'n dosis van FLEXOR KAPSULES te neem Neem die oorgeslaande dosis sodra jy onthou daarvan. Moet nie 'n dubbele dosis neem om op te maak vir individuele dosisse wat jy oorgeslaan of vergeet het nie. Uitwerking na behandeling met FLEXOR KAPSULES gestaak is Indien die dosis soos aangedui geneem word, sal daar geen nadelige newe-effekte wees wanneer die gebruik van hierdie medikasie gestaak word nie.
POSSIBLE SIDE EFFECTS FLEXOR CAPSULES can have side effects. Frequent side effects are nausea, heartburn, diarrhoea, constipation, bloating, burping and indigestion. Less frequent side effects are headache and heart palpitations. If any of the following happens, stop taking Flexor capsules and tell your doctor immediately or go to the casualty department at your nearest hospital: • Swelling of the hands, feet, ankles, face, lips, mouth or throat, which may cause difficulty in swallowing or breathing • Rash or itching • Fainting • Yellowing of the skin or eyes, also called jaundice. These are all very serious side effects. If you have any of them, you may have had a serious allergic reaction to Flexor Capsules. You may need urgent medical attention or hospitalisation. Tell your doctor immediately or go to the casualty department at your nearest hospital if you notice any of the following: • Spontaneous bleeding • Heart palpitations These are serious side effects. You may need urgent medical attention. Tell your doctor as soon as possible if you notice any of the following: nausea (feeling sick), heartburn, diarrhoea, constipation, bloating, burping, indigestion. If you notice any side effects not mentioned in this leaflet, please inform your doctor or pharmacist.	MOONTLIKE NEWE-EFFEKTE FLEXOR KAPSULES mag newe-effekte hê. Algemene newe effekte soos naarheid, sooibrand, diarree, hardlywigheid, opgeblaasheid, winde opbreek en spysvertering. Minder algemene newe-effekte is hoofpyn, lomerigheid en hartkloppings. Indien enige van die volgende gebeur staak Flexor Kapsules en raadpleeg jou dokter onmiddellik of gaan na die ongevalle afdeling van jou naaste hospitaal: • Swelling van die hande, voete, enkels, gesig, lippe, mond of keel wat die vermoë om te sluk of asem te haal bemoeilik. • Uitslag of jeukerigheid. • Floute. • Vergeling van die vel of oë, ook genaamd geelsug. Hierdie is ernstige newe-effekte. As jy enige van die simptome ervaar, kan jy 'n ernstige allergiese reaksie vir Flexor kapsules hê. Jy mag dan mediese sorg of hospitalisasie benodig. Vertel jou dokter onmiddellik of gaan na die ongevalle-afdeling van jou naaste hospitaal as jy enige van die volgende simptome het: • Spontane bloeding • Hartkloppings Hierdie is ernstige newe-effekte. Jy mag dringende mediese behandeling benodig. Vertel jou dokter onmiddellik as jy enige van die volgende agterkom: Naarheid, sooibrand, diarree, hardlywigheid, opgeblaasheid, winde opbreek, slegte spysvertering. As jy enige newe-effekte ervaar wat nie in die blaadjie genoem is nie, laat weet asseblief jou dokter of apteker.
STORING AND DISPOSING OF FLEXOR CAPSULES Store at or below 25 °C Protect from light and moisture. Store in original packaging. Do not store in a bathroom. Do not use after the expiry date printed on the bottle or box. Return all unused medicine to your pharmacist. Do not dispose of unused medicine in drains or sewage systems (e.g. toilets). STORE ALL MEDICINES OUT OF THE REACH OF CHILDREN.	HOE OM FLEXOR KAPSULES TE STOOR EN DAARVAN ONTSLAE TE RAAK Stoor onder 25 °C Beskerm teen lig en vog. Stoor / Bewaar in die oorspronklike verpakking. Moet nie in 'n badkamer stoor nie. Moet nie na die vervaldatum op die bottel of karton gebruik nie. Neem alle ongebruikte medikasie na jou apteker. Moet nie ongebruikte medikasie in dryne of rioolstelsels (bv. toilette) weggooi nie. STOOR ALLE MEDIKASIE BUITE BEREIK VAN KINDERS.
PRESENTATION OF FLEXOR CAPSULES The capsules are packed in a white plastic screw-top container containing 60 and 120 capsules.	AANBIEDING VAN FLEXOR KAPSULES Die kapsules is in n wit plastiek skroefdothouer wat 60 en 120 kapsules bevat.
IDENTIFICATION OF FLEXOR CAPSULES Off-white, 2 part capsule.	IDENTIFIKASIE VAN FLEXOR KAPSULES Afwit, twee deel kapsule.
REGISTRATION NUMBER Will be allocated by Council upon registration.	REGISTRASIENOMMER Sal toegeken word deur die Raad na afloop van registrasie.
NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION Keynote Health (Pty) Ltd 6 Heron Park 80 Corobrik Road Riverhorse Valley Durban 4017	NAAM EN BESIGHEIDSDRES VAN DIE HOUER VAN DIE REGISTRASIESERTIFIKAAT Keynote Health (Pty) Ltd 6 Heron Park 80 Corobrik Road Riverhorse Valley Durban 4017
DATE OF PUBLICATION March 2022	DATUM VAN PUBLIKASIE Maart 2022
This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.	Hierdie ongeregistreerde medisyne was nie geëvalueer deur die SAHPRA vir kwaliteit, veiligheid of gebruik intensies.